



SIMPLY WHITER TEETH™

*"Hey Doc,
I know you told me there might
be a little sensitivity, but nobody
told me whitening would hurt
like heck!*

*I can't do this...
I just want my money back."*

SOLVING TEETH WHITENING SENSITIVITY

By
Rod Kurthy
D.M.D.



Illustrations By Dr. Rod Kurthy

A KÖR Whitening
Science Paper

INTRODUCTION

Sensitivity (and sometimes downright pain) has been by far the most common negative side effect of teeth whitening – including both at-home and in-office whitening.¹⁻⁷ Millions of people have refrained from or discontinued whitening simply because of the discomfort.¹⁻⁷ As dentists, we hear this from people wherever we go, and especially in our dental practices.

Some attempts at eliminating sensitivity have resulted in whitening systems that didn't get teeth white. Some have resulted in whitening system claims of no sensitivity, when in fact sensitivity was not at all reduced.⁸⁻¹²

Recommended methods of preventing and treating whitening sensitivity abound, yet when dentists try these methods, often they find little or no benefit.⁹⁻¹² The causes of whitening sensitivity seem to be misunderstood by most, and methods of treating sensitivity are equally misunderstood.

CAUSES OF WHITENING SENSITIVITY

Dentists providing whitening services have found two distinct types of whitening sensitivity:

1) Typical Dentinal Hypersensitivity

Patients feel generalized discomfort of the teeth. Pulpals also overreact to various stimuli such as cold and teeth brushing.

2) Zingers

Zingers are those instantaneous sharp “electric shocks” that shoot down the length of anterior teeth, feeling like a lightning bolt, nearly bringing patients to their knees. They last for a couple seconds and are gone. The causes of Typical Dentinal Hypersensitivity are at least partially understood, however there is still speculation and controversy regarding the causes of zingers. We will present Dr. Rod Kurthy's hypothesis regarding zingers, which seems to have gained considerable attention by the dental community of late.

1. DENTINAL HYPERSENSITIVITY (DH)

DH is sometimes associated with genuine pathologic conditions such as caries, occlusal trauma, cracks in teeth, leakage under faulty restorations, etc. However our concern in this paper is dentinal hypersensitivity due to teeth whitening.

The Hydrodynamic Theory of Dentinal Hypersensitivity

The predominant theory of DH states that pulpal sensitivity is mediated by a “hydrodynamic mechanism”.^{1,13-17} A stimulus (thermal, mechanical, evaporative or osmotic/chemical) applied to dentin can increase the flow of dentinal tubular fluid within the tubules (either inward or outward).^{1,13-15,18,19} Flow of tubular fluid mechanically creates pressure or tension on the pulp, resulting in deformation of the cell membranes of A-delta nerve endings, which are mainly located at the periphery of the pulp at the inner surface of the dentin.^{1,14,20-24} (Fig. 2)

The term “hydraulic conductance” refers to the ease with which fluid flows through dentinal tubules.^{25,26} Obviously, the easier dentinal tubular fluid flows, the more and more often it flows, resulting in acute dentinal hypersensitivity.^{25,26}

Ideally, dentinal tubules should routinely have naturally-formed smear plugs blocking or plugging the entrance to the tubules,²⁵ thereby effectively reducing the tubular flow (conductance). (Fig. 1) When dentinal tubule smear plugs are lost, the hydraulic conductance increases a whopping 32-fold,^{25,27} resulting in potential for intense dentinal hypersensitivity. (Fig. 2)




Fig. 1 – Smear plugs are present in dentinal tubular orifices. The stimulus applied to dentinal surface has no effect on tubular fluid flow because of smear plugs. The pulp is protected and unaffected.




Fig. 2 – Smear plugs have been lost, allowing the stimulus applied to the dentinal surface to create outward tubular fluid flow, away from the pulp. Odontoblast processes are sucked deeper into the dentinal tubules, causing deformation of the odontoblasts and surrounding A-Delta Nerves, resulting in pain stimulation and inflammation.

Effects of an Acidic Diet

It is a fact that different people have different diameters of dentinal tubules^{25,28} and different degrees of flare of the orifices of those tubules.²⁵ Genetically, some have dentinal tubules twice the diameter of others.^{25,28} Acids not only decompose smear plugs and enlarge the inner diameter of the dentinal tubule, but they cause the orifice of the dentinal tubule to flare like a trombone.^{25,27} (Fig. 3 & 4)




Fig. 3 – Smear plugs are present in dentinal tubule orifices. Tubules have not been enlarged by acid challenges.




Fig. 4– Chronic acid challenges from acidic fruits, juices, sodas, sports drinks, acidic wine, etc., have removed the smear plugs, enlarged the inner diameter of the dentinal tubules and flared the orifices of the tubules.

Low salivary pH and an acidic diet (frequent consumption of fruits and fruit drinks, sodas, sports drinks, acidic wine, etc.) have long-term negative effects on dentinal tubules by enlarging the diameter and flaring the orifice of dentinal tubules.^{1,25,27} (Fig. 4) The larger the diameter of a tubule is, and the more the orifice flares, the easier it is to dislodge the tubular plug, and the more difficult it is to re-plug the dentinal tubule.²⁵

We have long known that people who eat significant amounts of acidic foods have more sensitivity.^{1,25,27} It is no surprise that statistically, patients with dentinal hypersensitivity tend to have dentinal tubules that are twice the diameter^{25,28} and often have very flared orifices compared to those who do not experience hypersensitivity.²⁵

Peroxides Dislodge Smear Plugs

The aggressive “oxygenation” phase of any whitening system results in the physical removal of smear plugs. (Fig. 5) Without smear plugs, hydraulic conductance of dentinal tubular fluid goes up 32 fold,^{25,27} and whitening gels can now intimately contact tubular fluids, resulting in a strong osmotic “pull” on the tubular fluid.



Fig. 5 – The aggressive “oxygenation” of the whitening gel is dislodging smear plugs from the orifices of dentinal tubules. The higher osmolarity whitening gel is osmotically drawing tubular fluid out of the tubules, initiating the hydrodynamic mechanism and creating sensitivity and discomfort.



Fig. 6 – The aggressive “oxygenation” of the whitening gel has removed smear plugs from the orifices of dentinal tubules. The higher osmolarity whitening gel is osmotically drawing tubular fluid out of the tubules, causing sensitivity and discomfort initiated by the hydrodynamic mechanism.

Whitening Gels Create an Osmotic Gradient

All whitening gels are hypertonic, with osmolalities varying from 4,900 mOsm/kg to 55,000 mOsm/kg, compared to only 290 mOsm/kg of dentinal tubular fluid.^{1,16} This means that whitening gels range from 17 to 190 times higher osmolality than dentinal tubular fluid.

The greater the osmolality of the whitening gel, the stronger the osmotic gradient between the gel and tubular fluid, the stronger the osmotic “pull” on the dentinal tubular fluid,^{14,16,17,29-31} and the more discomfort is felt by the patient.^{1,14,29,32,33} (Fig. 6)

Anhydrous and acidic pH whitening gels are more chemically stable,¹⁶ however both anhydrous and acidic gels have osmolalities that are up to eleven times greater than 100% aqueous and neutral pH gels.^{14,17,29-31} To avoid the costs of constant refrigeration, it is common for whitening product companies to use anhydrous gels and add acidifiers such as phosphoric acid to lengthen the shelf life.¹⁶

The more acidic and the more anhydrous the whitening gel is, the stronger the osmotic gradient is,^{14,29-31} the more forceful is the outward flow within the tubule,^{14,16,29-31} and therefore the more acute the discomfort may be for the patient.^{1,14,16,17,29,32,33}

Furthermore, acid in whitening gels more aggressively removes existing smear plugs within dentinal tubules, fostering more tubular fluid flow and even more sensitivity.^{1,27}

Peroxide Becomes More Acidic as it Decomposes

As peroxide decomposes, in addition to the formation of oxygen, oxygen ions and radicals; hydrogen ions are thrown off.³⁴ Hydrogen ions create acidity (the designation pH refers to “potential of Hydrogen”, and is a measure of the concentration of hydrogen ions).³⁴ This process can quickly cause an initially neutral gel to become acidic even down to a pH 3, further causing the problems noted above.

2. ZINGERS

During and/or soon after whitening, the patient may experience what feels like a sharp, immediate, intense “lightening bolt”, right down the length of an individual tooth. The patient may even describe this pain as a “crackling” electric shock. This tends to be spontaneous, for no apparent reason. It feels as if it starts at the incisal edge, and extends fully into the tooth.

Zingers usually occur in the smaller teeth – most often incisors and cuspids. They tend to occur in the same few teeth, over and over.

Peroxide Enters the Pulp

It has been shown that molecular hydrogen peroxide (H_2O_2) enters the pulp tissue during and after whitening.^{1,35-45} (Fig. 7) Contrary to what many of us were taught years ago, enamel is not nearly impervious.¹ In fact, it is now considered a semi-permeable membrane.¹ All teeth have “little highways” through tooth structure that allow low molecular weight hydrogen peroxide to enter the pulp.^{1,35-45} (Fig. 7) These aberrations (“little highways”) are formed naturally (enamel lamellae) or are acquired (cracks and craze lines).

Our Bodies Manufacture Hydrogen Peroxide

Our own bodies constantly produce large amounts of hydrogen peroxide and other oxidative chemicals, primarily in the mitochondria, during oxygen metabolism.⁴⁶⁻⁵⁰ The average adult human body produces about 650mg of hydrogen peroxide per day via oxygen metabolism.⁵¹ Two whitening trays with reservoirs, with 16% carbamide peroxide contain a total of only about 6.5mg of peroxide.⁵¹ This means that our own bodies manufacture approximately 100 times more peroxide every day than the amount that is put into upper and lower whitening trays.

If this large amount of daily endogenous hydrogen peroxide were allowed to continuously break down to strongly reactive oxygen species (such as free radicals) throughout our bodies, it would wreak havoc on our tissues.^{51,52} To protect against constant radical formation from the breakdown of peroxide, the body manufactures protective anti-oxidant enzymes including Catalase,⁵³⁻⁵⁵ superoxide dismutase, hemoxygenase-1 and glutathione peroxidase.^{46,52,56,57}

In the presence of these anti-oxidant enzymes, hydrogen peroxide is forced to break down to only water and molecular oxygen, instead of ions and radicals.⁵² This is precisely why hydrogen peroxide that enters the tooth pulp does not cause permanent damage.^{35,46,47,51-60}




Fig. 7 – Molecular hydrogen peroxide may enter the pulp during and after whitening through developmental or acquired aberrations through the enamel and dentin. The anti-oxidant enzyme Catalase present in the pulp breaks down the hydrogen peroxide to water and molecular oxygen. The formation of oxygen creates an instantaneous bubble in the pulp immediately increasing the intrapulpal pressure, aggressively distorting and stimulating the neurons of the pulp, creating a sharp painful “zinger” that lasts only seconds before the intrapulpal pressure dissipates.

Both gingival crevicular fluid and saliva contain high levels of peroxidase.^{16,48-50,61} To illustrate the effectiveness of peroxidase, salivary peroxidase alone can decompose 29mg of peroxide (4.5 times more than found in two whitening trays) per minute.⁵¹ There is therefore no concern about swallowing small amounts of peroxide gels.⁵¹

Catalase, which is always found in the dental pulp,^{35,53-55,62,63} is a very efficient enzyme molecule, and each Catalase molecule is capable of instantly breaking down several million molecules of hydrogen peroxide to water and molecular oxygen.⁶⁴

A common example of the effects of Catalase and peroxidase that most have observed is the bubbling seen when liquid hydrogen peroxide is poured into a cut. Hydrogen peroxide bubbles immediately upon contact with exposed bodily tissues and fluids. This reaction is due to the anti-oxidant enzymes such as Catalase⁵⁵⁻⁵⁷ and peroxidase within tissues and tissue fluids, which decompose the hydrogen peroxide to water and oxygen on contact.

DR. ROD KURTHY'S HYPOTHESIS OF ZINGER-TYPE WHITENING SENSITIVITY ETIOLOGY: (Fig. 7)

FACT: Molecular hydrogen peroxide (H_2O_2) enters the pulp during and after whitening through developmental or acquired aberrations in enamel and dentin.^{1,35-45}

FACT: Catalase is always found in the pulp.^{35,53-57,62,63}

FACT: When hydrogen peroxide comes into contact with Catalase, the hydrogen peroxide is instantly decomposed to water and oxygen gas.^{35,52,55-57}

HYPOTHESIS:

Because the pulp is housed in a rigid chamber inside the tooth, incapable of expanding, the instantaneous expansion of oxygen bubbles creates a very significant spike in intrapulpal pressure.¹⁴ This instantaneous spike in pressure deforms the membranes of virtually all pulpal neurons simultaneously, resulting in instant, intense pain throughout the entire pulp, feeling like a sharp electrical shock.¹⁴ (Fig. 7) The pressure quickly dissipates and equalizes, the deformation of the pulpal neurons ceases, and pain is gone. This hypothetical model perfectly describes the "zinger" type event.

PULPAL INFLAMMATION CAUSED BY BLEACHING LIGHTS AND LASERS

Whitening lights and lasers have been found to have no positive effect on teeth whitening results.⁶⁵⁻⁸⁶ When combined with higher concentration hydrogen peroxide, whitening lights and lasers were found to enhance pain^{16,17,73,76,77,84} and Substance P formation within the pulp, resulting in significantly higher pulpal inflammation and pain^{14,87} than when no lights or lasers were used.⁸⁸ Substance P is a neuropeptide released by pain transmitting neurons to communicate with each other. Its function is to cause pain and inflammation.^{14,87,88}

For a full explanation of the science related to whitening lights and lasers, including their significantly negative effect on the dental pulp, see the KöR Science Paper, *The Myth of Whitening Lights and Lasers*, by the same author.

CURRENT DESENSITIZING METHODS

There are two general categories of desensitizing product action:

1) Occlusion of dentinal tubules

By occluding (plugging) dentinal tubules, movement of intratubular fluid flow is prevented, and sensitivity is therefore prevented or ceased^{1,14,89} [treating the cause of sensitivity]. (Fig. 1)

2) Neuronal suppression

Chemical effect on pulpal neurons reducing the ability of pulpal neurons to fire.¹ [treating the symptoms of/masking sensitivity].

UNSUCCESSFUL ATTEMPTS OF REDUCING/ELIMINATING WHITENING SENSITIVITY

Modifying bleaching gels

In the past, several companies have altered whitening gels with intent to reduce sensitivity. What dentists have found is either:

- 1) these gels did reduce sensitivity, but also did not whiten teeth well
- 2) the claims of reduced sensitivity were false.⁸

Fluorides

Stannous and sodium fluoride combine with salivary calcium to create a precipitation of insoluble calcium fluoride within dentinal tubules.⁹⁰ The process of occlusion of tubules via use of fluoride requires extended treatment times and often never fully occludes tubules or cures sensitivity completely. Most dentists have not had remarkable results because of the very slow process of precipitation.

Use of fluorides in trays, as well as brushing with prescription strength fluoride for several weeks prior to whitening, during whitening, and after whitening has had some level of success, although rather minimal and unpredictable.^{91,92}

ACP

Amorphous Calcium Phosphate has primarily been discussed regarding remineralization of enamel, not dentin. The growth of inorganic hydroxyapatite within the fairly organic matrix of dentin is highly questionable.^{12,93,94}

Even if ACP were to actually promote mineralized formations within dentinal tubules, this process would not be immediate. The action of ACP in enamel is more of a “growth of hydroxyapatite” instead of a rapid occlusion of tubules.

Use of ACP has not been met with wide-spread reports of desensitizing success with whitening.^{12,93,94}

Potassium Nitrate

Potassium nitrate does not occlude tubules and does not reduce tubular flow.⁹⁵ It is theorized that it may reduce nerve excitability (inhibit re-polarization of pulpal neurons),¹ however the efficacy of potassium nitrate, having been around for decades, has not been strongly supported in the literature.^{1,9,91,92,95-99} To have an effect, potassium nitrate must migrate through dentinal tubules into the pulp,¹ which takes extended time.

Potassium nitrate has been unpredictable, seemingly effective on some, partially effective on some, and ineffective on some.^{1,99} Though potassium nitrate may reduce sensitivity in some individuals, it simply masks pulpal inflammation and does not eliminate it.

Addition of Desensitizers to Whitening Gels

Some whitening gels are manufactured with fluoride, potassium nitrate and/or ACP mixed into the whitening gels.^{9,10,12} Studies have not shown reduction of sensitivity from the addition of these substances into whitening gels.^{8,12}

Precipitation of calcium fluoride requires access to salivary calcium, which is excluded by whitening gel. Any effect by fluoride and ACP is a slow “growth” type process, which is interrupted by the aggressive chemistry and oxygenation process, as well as the outward flow of dentinal tubular fluid during whitening.

Potassium nitrate must migrate through the dentinal tubule to the pulp to have any positive benefit.¹ Whitening gels create an osmotic gradient resulting in outward flow within the dentinal tubules AWAY from the pulp.^{1,14,29-31} For potassium nitrate to reach the pulp during whitening, it would have to move through the tubule against the outward flow (like trying to swim upstream).^{14,29-31} Potassium nitrate mixed within the chemistry of whitening gel has been shown to be ineffective.^{8,9}

The chemical and aggressive oxygenation environment during whitening, as well as the outward flow of dentinal tubular fluid, is not conducive to the intended results of desensitizers of any type. Use of desensitizers is effective only before and/or after whitening, but not when mixed into the whitening gel itself.^{8,11,12}

SOLUTIONS TO WHITENING SENSITIVITY

Solutions to teeth whitening sensitivity include the following:

1. Creation of whitening gels with the lowest osmolality possible, to reduce the osmotic gradient between whitening gel and dentinal tubular fluid, thereby reducing intratubular flow and sensitivity.^{1,14,15,29-31}
 - a) Whitening gels should not only be aqueous,¹⁶ but 100% aqueous. To extend shelf life and gel stability during storage, constant refrigeration must be used instead of the use of anhydrous gels.
 - b) Whitening gels should be entirely neutral or even slightly alkaline.^{16,27} To extend shelf life and gel stability during storage, constant refrigeration must be used instead of adding acidifiers like phosphoric acid to whitening gels.
2. Prevention of dentinal hypersensitivity during whitening with rapid, aggressive occlusion of dentinal tubules, before and after each whitening activity instead of treating symptoms after they occur. Reinforcement of tubular smear plugs immediately prior to whitening, and rapid, aggressive replacement of any smear plugs lost during the oxygenation of whitening immediately after each whitening session results in predictability of success.^{14,25,89}
3. Prevention of zinger type whitening sensitivity with rapid, aggressive occlusion of enamel and dentin aberrations before, and after each whitening activity.
4. Use of desensitizer before and/or after whitening, but not mixed in with the whitening gel itself.⁸⁻¹²
5. Avoidance of whitening lights or lasers. Numerous studies have proven the ineffectiveness of lights and lasers.⁶⁵⁻⁷¹ With the ability to predictably accelerate whitening gels via pH and chemical acceleration,^{69,70} there is no need to consider the use of potentially harmful whitening lights or lasers.^{14,16,17,87,88}
6. Use of buffering agents to stabilize the neutral pH of whitening gels during decomposition in the mouth, preventing the natural tendency of peroxides to rapidly become acidic.

KÖR WHITENING SENSITIVITY SOLUTIONS

Evolve Dental Technologies is the first teeth whitening company to refrigerate a full line of whitening gels from the instant of manufacture until received cold by the dental office. Stabilization via refrigeration allows the whitening gels to be formulated with the lowest osmolality possible by using 100% aqueous gels and no acid content whatsoever.

The obvious benefit is the lowest possible osmotic gradient between KöR Whitening gels and the dentinal tubular fluid, the least amount of intratubular fluid flow, and therefore the least potential for sensitivity.^{1,14,15,29,32,33}

Dr. Kurthy's research indicates that rapid profound closure of dentinal tubules, immediately before and after whitening procedures, results in the most predictable whitening sensitivity control possible. (Figs. 11a & 11b)

OXALATE BASED DESENSITIZING

KöR Complete™ Desensitizer is a unique potassium oxalate desensitizer that strongly bonds to the highly mineralized enamel (96% hydroxyapatite) and peritubular dentin (92-93% hydroxyapatite) to provide immediate and profound closure of open dentinal tubules with no reduction in whitening effectiveness whatsoever. (Figs. 8,10a,10b,11a, & 11b)

Unlike typical oxalate desensitizing products that simply "clog up" the open orifices of dentinal tubules, KöR Complete reacts with and strongly bonds to the hydroxyapatite molecule in tooth structure via a chelation reaction, forming a durable compound of calcium oxalate salt crystals fused to peritubular hydroxyapatite 7-12µ deep within dentinal tubules. (Figs. 11a & 11b)

KöR desensitizers are used in the dental practice immediately before and after any in-office whitening session. They are not mixed in with the whitening gels. Desensitizer is also included with all at-home whitening kits, dispensed to patients to use daily after each at-home whitening to re-plug any tubules that may have been opened by the oxygenation process of whitening.¹⁰⁰ (Figs. 5,10a,10b,11a & 11b)

Fig. 8 Mineral (Hydroxyapatite) Content Percentages of Dental Hard Tissues

Enamel	≈ 96% hydroxyapatite (no collagen fibers or other proteins) ¹⁰¹
Peritubular dentin	≈ 92-93% hydroxyapatite (no collagen fibers or other proteins) ¹⁰¹
Intertubular dentin	≈ 70% hydroxyapatite (collagen fibers and other proteins found throughout) ¹⁰¹




Fig. 9 – SEM of dentinal surface. Note the highly mineralized peritubular dentin (≈ 92-93%hydroxyapatite with no collagen fibers or other proteins) surrounding open dentinal tubules.

Xu C, Wang Y. Chemical composition and structure of peritubular and intertubular human dentine revisited. Arch Oral Biol. 2012 Apr;57(4):383-91

University of Missouri - Kansas City School of Dentistry, Kansas City, Missouri



Fig. 10a – SEM photograph showing opened dentinal tubular orifice (loss of smear plug). Hydraulic conductance is 32 times greater than occluded tubules; tubular fluids are exposed to various stimuli capable of creating tubular fluid flow, resulting in tooth sensitivity and pain.




Fig. 10b – SEM photograph of open dentinal tubule – note peritubular dentin and intertubular dentin.



Fig. 11a – SEM photograph showing dental tubules closed and sealed with KöR Complete Desensitizer.




Fig. 11b – Same SEM photograph shown above.

Note peritubular dentin, intertubular dentin and calcium oxalate from KöR Complete Desensitizer sealing tubules.

KöR Complete has also been shown to bond to and within the natural enamel lamellae (developmental “cracks” through enamel, sometimes extending into dentin). Given the known ability of KöR Complete to strongly fuse to and seal enamel lamellae, it is believed that it also fuses to and within acquired craze lines and cracks in enamel.

With its ability to fuse to hydroxyapatite of enamel and dentin, KöR Complete forms a microscopically thin layer of calcium oxalate crystals over all tooth structure. Calcium oxalate crystals are snow-white, and therefore may even slightly increase the whiteness achieved during whitening procedures.

In addition to use with whitening procedures, KöR Complete Desensitizer has been shown to be a uniquely profound and durable desensitizer for virtually any clinical requirement, including treatment of generalized and localized tooth sensitivity; post periodontal surgery and hygiene; and under temporary/provisional restorations.

KÖR® DUAL-ACCELERATED, TRI-BARREL™ HYDREMIDE® PEROXIDE

With the use of a revolutionary Dual-Accelerated, Tri-Barrel™ Hydremide® Peroxide technology, the KÖR in-office whitening gels are chemically accelerated in two distinctly different ways, maximizing the effectiveness of the gels without the discomfort and potentially negative pulpal effects of a whitening light or laser.

This Tri-Barrel approach also enables the use of buffering agents that not only maintain a constant neutral pH throughout the whitening procedure, but also maintain the rapid effectiveness of the gels throughout the procedure. Remember, acidic pH stabilizes the whitening gel, and therefore slows its effectiveness. By maintaining a neutral pH throughout the breakdown of peroxide on the teeth, the whitening gel instability is maintained and whitening continues to progress quickly throughout the procedure.

There is currently no effective whitening system that can claim “no sensitivity”, however with the meticulous scientific approach discussed in this paper, both the incidence and severity of whitening related sensitivity has been greatly reduced by the KÖR desensitizing approach, and in the majority of cases virtually eliminated.

REFERENCES

- Pashley DH, Tay FR, Haywood VB, Collins MA, Drisko CL. Dentin Hypersensitivity: Consensus-Based Recommendations for the Diagnosis & Management of Dentin Hypersensitivity. *Inside Dentistry* (suppl). 2008 Oct; 4(9) (special issue).
- Haywood VB. *Tooth Whitening: Indications and outcomes of Nightguard Vital Bleaching*. Hanover Park, IL: Quintessence Publishing Company Inc; 2007.
- Browning WD, Blalock JS, Frazier KB, et al. Duration and timing of sensitivity related to bleaching. *J Esthet Restor Dent*. 2007;19(5):256-264.
- Settembrini L, Gultz J, Kaim J, Scherer W. A technique for bleaching nonvital teeth: inside/outside bleaching. *J Am Dent Assoc*. 1997;128(9):1283-1284.
- Swift EJ. Critical Appraisal: At-home bleaching: pulpal effects and tooth sensitivity issues, Part 1. *J Esthet Restor Dent*. 2006;18(4):225-228.
- Haywood VB, Cordero R, Wright K, et al. Brushing with a potassium nitrate dentifrice to reduce bleaching sensitivity. *J Clin Dent*. 2005;16(1):17-22.
- Haywood VB, Caughman WF, Frazier KB, et al. Tray delivery of potassium nitrate-fluoride to reduce bleaching sensitivity. *Quintessence Int*. 2001;32(2):105-109.
- Christensen GJ. At-Home Tooth Bleaching, State-Of-Art 2001. *Clinical Research Associates (CRA) Newsletter*. 2001 Feb;25(2):2-4
- Gallo JR, Burgess JO, Ripps AH, Bell MJ, Mercante DE, Davidson JM. Evaluation of 30% carbamide peroxide at-home bleaching gels with and without potassium nitrate – a pilot study. *Quintessence Int*. 2009 Apr;40(4):1-6.
- Tschoppe P, Neumann K, Mueller J, Kielbassa AM. Effect of fluoridated bleaching gels on the remineralization of predemineralized bovine enamel in vitro. *J Dent*. 2009 Feb;37(2):156-62. Epub 2008 Dec 11.
- Matis BA, Cochran MA, Eckert GJ, Matis JI. In vivo study of two carbamide peroxide gels with different desensitizing agents. *Oper Dent*. 2007 Nov-Dec;32(6):549-55.
- Giniger M, Spaid M, MacDonald J, Felix H. A 180-day clinical investigation of the tooth whitening efficacy of a bleaching gel with added amorphous calcium phosphate. *J Clin Dent*. 2005;16(1):11-6.
- Brännström M. A hydrodynamic mechanism in the transmission of pain produced stimuli through the dentine. In: *Sensory Mechanisms in Dentine*. Anderson DJ, ed. pp 73-79. Pergamon Press. London, 1963.
- Abd-Elmeguid A, Yu DC. Dental Pulp Neurophysiology: Part 1. Clinical and Diagnostic Implications. *JADC*. 2009;75(1):55.
- Braennstroem M, Astroem A. A study on the mechanism of pain elicited from the dentin. *J Dent Res*. 1964;43:619–25.
- Margeas RC. New advances in tooth whitening and dental cleaning technology. The Academy of Dental Therapeutics and Stomatology Dental Continuing Education Peer-Reviewed Web site. Accessed 2010;March.
- Papathanasiou A, et al. Clinical evaluation of a 35% hydrogen peroxide in-office whitening system. *Comp*. 2002;23:335–346.
- Marvin K. Bright, White, and Sensitive: An Overview of Tooth Whitening and Dentin Hypersensitivity. *Dentistry Today.com*. 2009 Sept.
- Drisko CH. Dentine hypersensitivity: dental hygiene and periodontal considerations. *Int Dent J*. 2002;52:385-393.
- Bergenholtz G, Hörsted-Bindslev P, Reit C. *Textbook of endodontology*. Oxford, UK: Blackwell Munksgaard; 2003.
- Nanci A, Ten Cate AR. *Ten Cate's oral histology: development, structure, and function*. 6th ed. St. Louis: Mosby; 2003.
- Andrew D, Matthews B. Displacement of the contents of dentinal tubules and sensory transduction in intradental nerves of the cat. *J Physiol*. 2000; 529(3):791–802.
- Narhi MV. Dentin sensitivity: a review. *J Biol Buccale*. 1985; 13(2):75–96.
- Narhi M, Jyvasjarvi E, Virtanen A, Huopaniemi T, Ngassapa D, Hirvonen, T. Role of intradental A- and C-type nerve fibres in dental pain mechanisms. *Proc Finn Dent Soc*. 1992; 88 Suppl 1:507–516.
- Absi EG, Addy M, Adams D. Dentine hypersensitivity. A study of the patency of dentinal tubules in sensitive and non-sensitive cervical dentine. *J Clin Periodontol*. 1987 May;14(5):280-4.
- Yoshiyama M, Noiri Y, Ozaki K, Uchida A, Ishikawa Y, Ishida H. Transmission electron microscopic characterization of hypersensitive human radicular dentin. *J Dent Res*. 1990;69:1293-7.
- Reeder OW Jr, Walton RE, Livingston MJ, Pashley DH. Dentin permeability: determinants of hydraulic conductance. *J Dent Res*. 1978 Feb;57(2):187-93.
- Stojis I, Petrovi L, Stojanac I, Drobnac M. Multi-factoriality of dentine hypersensitivity. *Medicinski Pregled*. 2008 Jul-Aug;61(7-8):359-63.
- Anderson DJ, Matthews B, Shelton LE. Variations in the sensitivity to osmotic stimulation of human dentine. *Arch Oral Biol*. 1967; 12(1):43–7.

REFERENCES (continued)

30. Narhi M, Kontturi-Narhi V, Hirvonen T, Ngassapa D. Neurophysiological mechanisms of dentin hypersensitivity. *Proc Finn Dent Soc.* 1992; 88 Suppl 1:15–22.
31. Narhi MV, Hirvonen T. The response of dog intradental nerves to hypertonic solutions of CaCl₂ and NaCl, and other stimuli, applied to exposed dentine. *Arch Oral Biol.* 1987;32(11):781–6.
32. Vongsavan N, Matthews B. The relationship between the discharge of interdental nerves and the rate of fluid flow through dentine in the cat. *Arch Oral Biol.* 2007 Feb;52(7):640–7.
33. Pashley DH. Sensitivity of dentin to chemical stimuli. *Endod Dent Traumatol.* 1986; 2(4):130–7.
34. Eary LE. Catalytic decomposition of hydrogen peroxide by ferric ion in dilute sulfuric acid solutions. *Metallurgical and Materials Transactions B.* 1985 June; 16(2):181–186.
35. Bowles WH, Burns H Jr. Catalase/peroxidase activity in dental pulp. *J Endod.* 1992 Nov;18(11):527–34.
36. Jorgensen MG, Carroll WB. Incidence of tooth sensitivity after home whitening treatment. *J Am Dent Assoc* 2002; 133: 1076–1082.
37. Pohjola RM, Browning WD, Hackman ST, Myers ML, Downey MC. Sensitivity and tooth whitening agents. *J Esthet Restor Dent* 2002; 14: 85–91.
38. Matis BA, Mousa HN, Cochran MA, Eckert GJ. Clinical evaluation of bleaching agents of different concentrations. *Quintessence Int* 2000; 31: 303–310.
39. Bowles WH, Ugwuneri Z. Pulp chamber penetration of hydrogen peroxide following vital bleaching procedures. *J Endodont* 1987; 13: 375–377.
40. Cooper J, Bokmeyer T, Bowles W. Penetration of the pulp chamber by bleaching agents. *J Endodont* 1992; 18: 315–317.
41. Thitinanthapan W, Satamanont P, Vongsavan N. In vitro penetration of the pulp chamber by three brands of carbamide peroxide. *Esthet Dent* 1999; 11: 259–264.
42. Slezak B, Santarpia P, Xu T, et al. Safety profile of a new liquid whitening gel. *Compend Contin Educ Dent* 2002; 23(Suppl 1): 4–11.
43. Pugh G, Zaidel L, Lin N, Stranick M, Bagley D. High levels of hydrogen peroxide in overnight tooth-whitening formulas: effects on enamel and pulp. *J Esthet Restor Dent* 2005; 17: 40–45.
44. Sulieman M. An overview of bleaching techniques: 3. In-surgery or power bleaching. *Dent Update* 2005; 32: 101–104, 107–108.
45. Sulieman M, Addy M, Macdonald E, Rees JS. A safety study in vitro for the effects of an in-office bleaching system on the integrity of enamel and dentine. *J Dent* 2004; 32: 581–590.
46. Anderson D, Chiego Jr. D, Glickman G, McCauley L. A clinical assessment of the effects of 10% carbamide peroxide gel on human pulp tissue. *Journal of Endodontics.* 1999;25(4):247–250.
47. Fugaro JO, Nordahl I, Fugaro OJ, et al. Pulp reaction to vital bleaching. *Oper Dent.* 2004;29(4):363–368.
48. Patel S, Pradeep A, Chowdhry S. Crevicular fluid levels of plasma glutathione peroxidase (eGPx) in periodontal health and disease. *Archives of Oral Biology.* 2009 Jun;54(6):543–8.
49. Jentsch H, Sievert Y, Göck R. Lactoferrin and other markers from gingival crevicular fluid and saliva before and after periodontal treatment. *Journal of Clinical Periodontology.* 2004 Jul;31(7):511–4.
50. Kaner D, Bernimoulin JP, Kleber BM, Heizmann WR, Friedmann A. Gingival crevicular fluid levels of calprotectin and myeloperoxidase during therapy for generalized aggressive periodontitis. *J Periodontal Research.* 2006 Apr;41(2):132–9.
51. Haywood V. Tray Bleaching Overview and Indications. 2009 Western Dental Regional Dental Convention. Phoenix, AZ; March 12–14, 2009:track 12, disk 1/5.
52. Enzyme-Catalase. <http://www.enzymeindia.com/Enzymes-Catalase.php>. Advanced Enzyme Technologies Ltd. Chino CA, USA.
53. Babior BM. Oxidants by phagocytes: agents of defense and destruction. *Blood.* 1984;64:959–966.
54. Forman HJ, Torres M. Redox signaling in macrophages. *Mol Aspects Med.* 2001;22:189–216.
55. Roos D, Weening RS, Wyss SR, Aebi HE. Protection of human neutrophils by endogenous catalase: studies with cells from catalase-deficient individuals. *J Clin Invest.* 1980;65:1515–1522.
56. Esposito P, Varvara G, Murmura G, Terlizzi A, Caputi S. Ability of healthy and inflamed human dental pulp to reduce hydrogen peroxide. *Eur J Oral Sci.* 2003 Oct;111(5):454–6.
57. Esposito P, Varvara G, Caputi S, Perinetti G. Catalase activity in human healthy and inflamed dental pulps. *Int Endod J.* 2003 Sep;36(9):599–603.
58. RIGOBELLO MP, FOLDA A, BALDOIN MC, SCUTARI G, BINDOLI A. Effect of auranofin on the mitochondrial generation of hydrogen peroxide. Role of thioredoxin reductase. *J Free radical research.* 2005;39(7):687–695.
59. LEE SJ, JIN Y, HYE YY, CHOI B, HYOUNG CK, OH YK, KIM HS, KIM WK. Cyclopirox protects mitochondria from hydrogen peroxide toxicity. *British J Pharm.* 2005;145(4):469–476.
60. Dionisi O, Galeotti T, Terranova T, Azzi A. Superoxide radicals and hydrogen peroxide formation in mitochondria from normal and neoplastic tissues. *Biochem Biophys Acta.* 1975 Oct 22;403(2):292–300
61. Wei PF, Ho KY, Ho YP, Wu YM, Yang YH, Tsai CC. The investigation of glutathione peroxidase, lactoferrin, myeloperoxidase and interleukin-1 in gingival crevicular fluid: implications for oxidative stress in human periodontal diseases. *J Periodontal Research.* 2004 Oct;39(5):287–293.
62. Davis WL, Jacoby BH, Craig KR, Wagner G, Harrison JW. Copper-zinc superoxide dismutase activity in normal and inflamed human dental pulp tissue. *J Endodont.* 1991;17:316–318.
63. Freeman BA, Crapo JD. Biology of disease: free radicals and tissue injury. *Lab Invest.* 1982;47:412–426.
64. Catalase. Wikipedia. 06 March 2010. <http://en.wikipedia.org/wiki/Catalase>
65. Bruzell EM, Johnsen B, Aalerud TN, Dahl JE, Christensen T. In vitro efficacy and risk for adverse effects of light-assisted tooth bleaching. *Photochem Photobiol Sci.* 2009 Mar;8(3):377–85.
66. Lima DA, Aguiar FH, Liporoni PC, Munin E, Ambrosano GM, Lovadino JR. In vitro evaluation of the effectiveness of bleaching agents activated by different light sources. *J Prosthodont.* 2009 Apr;18(3):249–54.
67. Christensen GJ. New Generation In-office Vital Tooth Bleaching, Part 2. *Clinical Research Associates (CRA) Newsletter.* 2003 March:27(3):1–3
68. Haywood, V. Masters of Esthetic Dentistry. *Journal of Esthetic and Restorative Dentistry.* 2003;15(3).
69. Hein DK, Ploeger BJ, Hartup JK, Wagstaff RS, Palmer TM, Hansen LD. In-office vital tooth bleaching--what do lights add? *Comp Contin Edu Dent.* 2003 Apr;24(4A):340–52.
70. Kugel G, Papatheanasiou A, Williams AJ 3rd, Anderson C, Ferreira S. Clinical evaluation of chemical and light-activated tooth whitening systems. *Compend Contin Educ Dent.* 2006 Jan;27(1):54–62.
71. Jones AH, Diaz-Arnold AM, Vargas MA, Cobb DS. Colorimetric assessment of laser and home bleaching techniques. *J Esthet Dent.* 1999;11(2):87–94.
72. Baroudi K, Hassan NA. The effect of light-activation sources on tooth bleaching. *Niger Med J.* 2014 Sep-Oct; 55(5): 363–368.
73. Kossatz S, Dalanhol AP, Cunha T, Loguercio A, Reis A. Effect of light activation on tooth sensitivity after in-office bleaching. *Oper Dent.* 2011 May-Jun;36(3):251–7. doi: 10.2341/10-289-C. Epub 2011 Jul 8.
74. Christensen G, Tooth Bleaching, State-of-Art '97. *Clinical Research Associates Newsletter* 1997;21(4).

REFERENCES (continued)

75. Strobl A, Gutknecht N, Franzen R, Hilgers RD, Lampert F, Meister J. Laser-assisted in-office bleaching using a neodymium:yttrium-aluminum-garnet laser: an in vivo study. *Lasers in Medical Science*. 2009; May.
76. Kugel G, Ferreira S, Sharma S, Barker ML, Gerlach RW. Clinical trial assessing light enhancement of in-office tooth whitening. *J Esthet Restor Dent*. 2009;21(5):336-47
77. Marson FC, Sensi LG, Vieira LC, Araujo E. Clinical evaluation of in-office dental bleaching treatments with and without the use of light-activation sources. *Oper Dent*. 2008;33:15-22.
78. Bernardon JK, Sartori N, Ballarin A, Perdigão J, Lopes GC, Baratieri LN. Clinical performance of vital bleaching techniques. *Oper Dent*. 2010 Jan-Feb; 35(1):3-10.
79. Hahn P, Schondelmaier N, Wolkewitz M, Altenburger MJ, Polydorou O. Efficacy of tooth bleaching with and without light activation and its effect on the pulp temperature: an in vitro study. *Odontology*. 2013 Jan; 101(1):67-74.
80. Nutter BJ, Sharif MO, Smith AB, Brunton PA. A clinical study comparing the efficacy of light activated in-surgery whitening versus in-surgery whitening without light activation. *J Dent*. 2013 Nov; 41 Suppl 5():e3-7.
81. Liang S, Sa Y, Jiang T, Ma X, Xing W, Wang Z, et al. In vitro evaluation of halogen light-activated vs chemically activated in-office bleaching systems. *Acta Odontol Scand*. 2013;71:1149-55.
82. Roberto AR, Jassé FF, Boaventura JMC, Martinez TC, Rastelli ANS, Oliveira OB Jr, Saad JRCS. Evaluation of tooth color after bleaching with and without light-activation. *Rev Odonto Cienc* 2011;26(3):247-252.
83. Kugel G, Ferreira S. The art and science of tooth whitening. *J Mass Dent Soc*. 2005 Winter;53(4):34-7.
84. He LB, Shao MY, Tan K, Xu X, Li JY. The effects of light on bleaching and tooth sensitivity during in-office vital bleaching: a systematic review and meta-analysis. *J Dent*. 2012 Aug;40(8):644-53.
85. de Almeida Farhat PB, Santos FA, Gomes JC, Gomes OM. Evaluation of the efficacy of LED-laser treatment and control of tooth sensitivity during in-office bleaching procedures. *Photomed Laser Surg*. 2014 Jul;32(7):422-6.
86. Alomari O, El Daraa E. A randomized clinical trial of in-office dental bleaching with or without light activation. *J Contemp Dent Pract*. 2010 Jan 1;11(1):E017-24.
87. Z Olgart L. Neural control of pulpal blood flow. *Crit Rev Oral Biol Med*. 1996;7(2):159-71.
88. Caviedes-Bucheli J, Ariza-García G, Restrepo-Méndez S, Ríos-Orsorio N, Lombana N, Muñoz HR. The effect of tooth bleaching on substance P expression in human dental pulp. *Journal of Endodontics*. 2008 Dec;34(12):1462-5.
89. Byers MR. Effects of inflammation on dental sensory nerves and vice versa. *Proc Finn Dent Soc*. 1992;88 Suppl 1:499-506.
90. Orchardson R, Gillam DG. Managing dentin hypersensitivity. *J Am Dent Assoc*. 2006;137(7):990-998.
91. Docimo R, Montesani L, Maturo P, Costacurta M, Bartolino M, Zhang YP, DeVizio W, Delgado E, Cummins D, Dibart S, Mateo LR. Comparing the efficacy in reducing dentin hypersensitivity of a new toothpaste containing 8.0% arginine, calcium carbonate, and 1450 ppm fluoride to a benchmark commercial desensitizing toothpaste containing 2% potassium ion: an eight week clinical study in Rome, Italy. *J Clin Dent*. 2009;20(4):137-43.
92. Nathoo S, Delgado E, Zhang YP, DeVizio W, Cummins D, Mateo LR. Comparing the efficacy in providing instant relief of dentin hypersensitivity of a new toothpaste containing 8.0% arginine, calcium carbonate, and 1450 ppm fluoride relative to a benchmark desensitizing toothpaste containing 2% potassium ion and 1450 ppm fluoride, and to a control toothpaste with 1450 ppm fluoride: a three-day clinical study in New Jersey, USA. *J Clin Dent*. 2009;20(4):123-30.
93. Fiocchi MF, Moretti AJ, Powers JM, Rives T. Treatment of root sensitivity after periodontal therapy. *Am J Dent*. 2007 Aug;20(4):217-20.
94. Yates R, Owens J, Jackson R, Newcombe RG, Addy M. A split-mouth placebo-controlled study to determine the effect of amorphous calcium phosphate in the treatment of dentine hypersensitivity. *J Clin Periodontol*. 1998 Aug;25(8):687-92.
95. Pereira R, Chava VK. Effects of a potassium nitrate mouthwash on dentinal tubules--a SEM analysis using the dentine disc model. *J Int Acad Periodontol*. 2002 Apr;4(2):44-8.
96. Poulsen S, Errboe M, Hovgaard O, Worthington HW. Potassium nitrate toothpaste for dentine hypersensitivity. *Cochrane Database Syst Rev*. 2001; (2):CD001476.
97. Poulsen S, Errboe M, Lescay Mevil Y, Glenny AM. Potassium containing toothpastes for dentine hypersensitivity. *Cochrane Database Syst Rev*. 2006 Jul 19;3:CD001476.
98. Kishore A, Mehrotra KK, Saimbi CS. Effectiveness of desensitizing agents. *J Endod*. 2002 Jan;28(1):34-5.
99. Orchardson R, Gillam DG. Managing dentin hypersensitivity. *J Am Dent Assoc*. 2006;137(7):990-998.
100. Sykes LM. Dentine hypersensitivity: a review of its aetiology, pathogenesis and management. *SADJ*. 2007 Mar;62(2):066-71.
101. Xu C, Wang Y. Chemical composition and structure of peritubular and intertubular human dentine revisited. *Arch Oral Biol*. 2012 Apr;57(4):383-91.



SIMPLY WHITER TEETH™

www.KoRWhitening.com
866-763-7753

5 Vanderbilt, Irvine, CA 92618